

ANNEX 3. DOSAGES FOR ARV DRUGS

Dosages of ARV drugs for adults and adolescents

Generic name	Dose
Nucleoside reverse-transcriptase inhibitors (NRTIs)	
Abacavir (ABC)	300 mg twice daily or 600 mg once daily
Emtricitabine (FTC)	200 mg once daily
Lamivudine (3TC)	150 mg twice daily or 300 mg once daily
Zidovudine (AZT)	300 mg twice daily
Nucleotide reverse-transcriptase inhibitors (NtRTIs)	
Tenofovir disoproxil fumarate (TDF)	300 mg once daily
Tenofovir alafenamide (TAF)	10-25 mg once daily ^a
Non-nucleoside reverse-transcriptase inhibitors (NNRTIs)	
Efavirenz (EFV)	400–600 mg once daily
Etravirine (ETV)	200 mg twice daily
Nevirapine (NVP)	200 mg once daily for 14 days followed by 200 mg twice daily
Proteases inhibitors (PIs)	
Atazanavir/ritonavir (ATV/r)	300 mg/100 mg once daily
Darunavir + ritonavir (DRV/r)	800 mg + 100 mg once daily or 600 mg + 100 mg twice daily
Lopinavir/ritonavir (LPV/r)	400 mg/100 mg twice daily
	Considerations for individuals receiving TB therapy In the presence of rifampicin, adjusted dose of LPV/r ("Double dose" LPV 800 mg/ + ritonavir 200 mg twice daily or "super boosted" with LPV 400 mg/ + ritonavir 100 mg twice daily plus additional doses of RTV 300 mg twice daily), with close monitoring. In the presence of rifabutin, no dose adjustment required. Rifapentine should not be used.
Integrase strand transfer inhibitors (integrase inhibitors)	
Dolutegravir (DTG)	50 mg once daily ^b
Raltegravir (RAL)	400 mg twice daily
	Considerations for individuals receiving TB therapy In the presence of rifampicin, adjusted dose of DTG (50 mg twice daily) and RAL (800 mg twice daily), with close monitoring. In the presence of rifabutin or rifapentine, no dose adjustment is required.

^a TAF 25 mg and TAF/FTC/DTG (TAF 25 mg, Emtricitabine 200 mg, Dolutegravir 50 mg fixed dose combination) can be used once daily in adolescents living with HIV weighing at least 25 kg.

^b DTG 50 mg and TLD (Tenofovir 300 mg, Lamivudine 300 mg, Dolutegravir 50 mg fixed dose combination) can be used once daily in adolescents living with HIV weighing at least 30 kg.

WEIGHT-BASED DOSING FOR ARV FORMULATIONS FOR INFANTS AND CHILDREN

Prescribing information and weight-based dosing of available ARV formulations for infants and children

This annex contains information on ARV drugs for which there are paediatric indications, formulations or sufficient information and evidence to provide guidance on prescribing and dosing in infants, children and adolescents less than 18 years of age. The work to develop and update simplified guidance on ARV drugs for use in children has been undertaken by WHO through the Paediatric Antiretroviral Working Group¹.

For simplification and ease of implementation, doses are expressed per weight-band rather than per kilogram or per square metre of body surface area. When this simplified weight-band dosing was developed, careful consideration was given to the expected body surface area of children from low- and middle-income countries in each weight band. The primary source of information for the guidance provided is the manufacturer's package insert. This was supplemented with data from other clinical studies as well as expert paediatric pharmacology consultations. For fixed-dose combinations, a dose-modelling tool (<http://www.who.int/hiv/paediatric/generictool/en/index.html>) was used to predict the dose delivered for each component drug against the recommended dosing schedule. In some cases, the dose for a component in a particular weight band may be somewhat above or below the target dose recommended by the manufacturer. This is inevitable given the limitations imposed by a fixed-dose combination, but care was taken to ensure that in no case would a child receive more than 25% above the maximum target dose or more than 5% below the minimum target dose. PK efficacy and safety studies have also confirmed the overall safety of this dosing approach. For simplification, ARV drugs no longer considered preferred or alternative options for children have been removed from the dosing guidance.

In the context of increasing implementation of HIV virological testing at birth, and the shift towards treating infants earlier in an effort to reduce early mortality, these guidelines include weight-based dosing for term infants aged <4 weeks, including those weighing less than 3 kg. However, there is limited experience with initiating treatment in neonates with HIV infection aged <2 weeks, and a paucity of PK data to fully inform accurate dosing for most drugs in neonates, who are undergoing rapid growth and maturation in renal and liver function. Limited PK data in preterm infants are available for AZT, NVP, and 3TC; there is considerable uncertainty of appropriate dosing for NVP, RAL and 3TC in preterm and low birth weight infants. In addition, LPV/r solution should not be given to preterm infants until they have reached 42 weeks gestational age, because of the risk of adverse effects that may occur in this population. The management of HIV treatment in preterm neonates is extremely challenging because of the lack of appropriate pharmacokinetic, safety, and dosing information as well as suitable formulations. Dosing for postnatal prophylaxis for infants exposed to HIV is not provided here but can be found [here](#).

In the 2018 ARV guidelines revision, integrase inhibitors had been included more prominently among the preferred regimens recommended by WHO. At the time of this guidelines update in 2019, DTG was only approved for children above 6 years and 15 kg in Europe² and above 30 kg in the United States³. The registration trial⁴ is anticipated to generate data for dosing DTG in children down to age 4 weeks by the end of 2019 in 5 mg increments, with potential regulatory approval in mid-2020. This dosing annex includes approved DTG dosing as well as simplified dosing based on PK and safety data from an ongoing multi-country study, which is also investigating the pharmacokinetics of DTG in TB co-treated children⁵. Findings from this trial led to a further revision of the DTG dosing in January 14th 2019, when the PAWG reviewed and discussed PK data for the 20-24.9 kg weight band. The PAWG members acknowledged the short follow up and limited clinical experience, but had no major concerns with adopting a simplified dosing for this weight band and agreed with recommending use of DTG 50 mg tablets from 20 kg. As the introduction of paediatric formulations of DTG will take time, programmes are encouraged to begin using DTG 50 mg in children weighing 20 kg or more while planning for the future use of DTG in younger paediatric populations while the simplified dosing is being confirmed and appropriate formulations are being developed.

RAL granules were also added with the goal of providing a suitable formulation to deliver RAL to neonates. Due to concerns about the complexity of administration of the granule formulation, the 25 mg chewable tablets as disperible tablets have been endorsed by the PAWG for infants and children older than 4 weeks of age and weighting at least 3 kg. This decision was largely based on in vitro data on solubility and bioequivalence between RAL tablets and granules⁶ as well as considering the limited availability of adequate alternative formulations for this age group.

In this 2019 update, paediatric dosing information for tenofovir alafenamide (TAF) and fixed dose combination containing TAF were also included for children weighing 25 kg or more. This aligns with dosing approved by US FDA⁷. Studies to investigate dosing below 25 kg are ongoing and more information will be made available as soon as approval is extended.

This dosing annex and the simplified dosing schedule will be regularly reviewed and updated as additional data and new formulations become available.

Antiretroviral drugs and formulations are available from several manufacturers, and available dosage strengths of tablets, capsules and liquid formulations may vary from the information provided here. Several optimal paediatric dosage forms are currently in development but have not yet received regulatory approval at the time of writing these updated guidelines. National programme managers should ensure that products planned for use have received stringent regulatory approval and are of appropriate quality and stability. For guidance on the quality assurance of medicines, see the WHO medicines web site (http://www.who.int/medicines/areas/quality_safety/quality_assurance/about/en/index.html) and the *Access to HIV/AIDS drugs and diagnostics of acceptable quality*, which is available and updated at <http://www.who.int/hiv/amds/selection/en/index.html>. The current list of WHO prequalified drugs is available at <http://apps.who.int/prequal>. For the current list of ARV drugs approved and tentatively approved by the United States Food and Drug Administration, see: <https://www.fda.gov/InternationalPrograms/PEPFAR/ucm119231.htm>. For the policy of the Global Fund to Fight AIDS, Tuberculosis and Malaria on procurement and quality assurance, see <https://www.theglobalfund.org/en/sourcing-management/quality-assurance/medicines/>

General principles

The principles followed in developing the WHO simplified tables include the following:

- Use of an age-appropriate fixed dose combination is preferred for any regimen if such a formulation is available.
- Oral liquid or syrup formulations should be avoided where possible. Dispersible tablets (or granules for oral solution) are the preferred solid oral dosage forms, since each tablet can be made into liquid at the point of use.
- If suitable dispersible FDC's are not available and oral liquids must be used, it is recommended that children be switched to a solid oral dosage form as soon as possible.
- While dosing neonates generally necessitates use of oral liquid formulations for administrating precise dosing, switching to solid oral dosage form as soon as possible is recommended.
- Where children have to use adult formulations, care must be taken to avoid under-dosing and overdosing. Use of scored tablets are preferred to ensure accurate dosing is provided, particularly if adult dosage forms are used. Splitting of unscored tablets should be avoided as uniform distribution of active drug product cannot be assured in tablet fragments.
- Some tablets such as LPV/r or ATV/r heat-stable tablets are made in a special embedded matrix formulation (a proprietary melt extrusion technology that stabilizes drug molecules that are normally heat labile) and should not be cut, split, dissolved, chewed or crushed, since bioavailability is significantly reduced when not swallowed whole.
- LPV/r is available in a 40 mg/10 mg pellet or granule formulation for infants and young children. However, children weighing 10 kg or more should be transitioned to LPV/r heat-stable tablets as soon as they are able to swallow tablets whole.
- At each clinic visit, children should be weighed and doses should be adjusted based on observed growth and change in body weight.
- Country programs should consider the national regulatory status and local availability status of specific dosage forms when developing national paediatric treatment recommendations.
- Research is ongoing for several antiretroviral medications to establish dosing guidance in neonates, infants and young children. The age indications for each drug mentioned in the drug pages are based on current evidence and will be updated as new recommendations become available.

Table 1 Simplified dosing of child-friendly fixed-dose solid formulations for twice-daily dosing in infants and children 4 weeks of age and older^a

Drug	Strength of paediatric tablets	Number of tablets by weight band morning and evening						Strength of adult tablet	Number of tablets by weight band
		3–5.9 kg	6–9.9 kg	10–13.9 kg	14–19.9 kg	20–24.9 kg	25–34.9 kg		
		AM	PM	AM	PM	AM	PM	AM	PM
AZT/3TC 60 mg/30 mg	Tablet (dispersible)	1	1	1.5	1.5	2	2	2.5	2.5
AZT/3TC/ NVP ^b	Tablet (dispersible) 60 mg/30 mg/50 mg	1	1	1.5	1.5	2	2	2.5	3
ABC/3TC 60 mg/30 mg ^c	Tablet (dispersible) 120/60 mg	0.5	0.5	0.5	1	1	1	1.5	1.5

^a For infants younger than 4 weeks of age refer to table 4 for more accurate dosing which is reduced due to the decreased ability to excrete and metabolize medications. For infants who are at least 4 weeks of age but less than 3 kg, immaturity of renal and hepatic pathways of elimination are less of a concern but uncertainty still exists on the appropriate dosing of ARVs in preterm and low birth weight infants.

^b Please note that this regimen and formulation is no longer recommended and should only be used in special circumstances where other age appropriate formulations are not available.

^c This formulation will be phased out of use over time and programs should transition to use of the 120 mg/60 mg dispersible scored tablets.

Table 2 Simplified dosing of child-friendly solid formulations for once-daily dosing in infants and children 4 weeks of age and older^a

Drug	Strength of paediatric tablet	Number of tablets or capsules by weight band once daily				Strength of adult tablet	Number of tablets or capsules by weight band once daily
		3–5.9 kg	6–9.9 kg	10–13.9 kg	14–19.9 kg		
EFV ^b	Tablet (scored) 200 mg	—	—	1	1.5	1.5	—
ABC/3TC	Tablet (dispersible) 60/30 mg	2	3	4	5	6	2
ABC/3TC	Tablet (dispersible) 120/60 mg	1	1.5	2	2.5	3	—
TAF ^c	Tablet 25 mg	—	—	—	—	—	—
ATV ^d	Capsules 100 mg	—	—	2	2	2	1
	Capsules 200 mg	—	—	1	1	1	1 ^e
DRV ^f	Tablet 600 mg	—	—	—	1	1	600 mg
	Tablet 150 mg	—	—	—	4	4	—
RTV ^g	Tablet 25 mg	—	—	—	4	4	100 mg
	Tablet 50 mg	—	—	—	2	2	—
DTG ^h	Film-coated Tablet 50 mg	—	—	—	1	1	50 mg

^a See table 4 for dosing recommendations for infants younger than 4 weeks old. Doses for this age group are reduced to account for the decreased ability to excrete and metabolize medications. For infants who are at least 4 weeks of age but less than 3 kg, immaturity of renal and hepatic pathways of elimination are less of a concern but uncertainty still exists on the appropriate dosing of ARVs in preterm and low birth weight infants.

^b EEFV is not recommended for children younger than 3 years and weighing less than 10 kg.

^c At the time of this update, TAF film coated tablets were approved for children above 6 years by FDA for use in un-boosted regimens such as with DTG. A fixed dose combination containing TAF/FTC/DTG (TAF 25 mg, Emtricitabine 200 mg, Dolutegravir 50 mg fixed dose combination) received tentative approval by US FDA and can be used once daily in children and adolescents living with HIV weighing at least 25 kg.

^aATV is only approved for use in children 3 months and older. ATV single strength capsules should be administered with RTV 100 mg for all weight bands 10 kg and above. ATV powder formulation has limited availability in LMIC, but enables administration of ATV to infants and children as young as 3 months. Infants and children 5-15 kg should be administered 200 mg of ATV powder (4 packets, 50 mg/packet) with 80 mg of RTV oral solution (1 ml). https://www.accessdata.fda.gov/drugsatfda_docs/label/2018/021567s042_206352s071b1.pdf

^bA 300 mg dose for 25-29.9 kg is recommended on the basis of findings from the PRINCE-2 study⁸

^cDRV in combination with RTV should be used, in children older than 3 years, once daily when this is used without previous exposure to PI. While approved dosing for 30-35kg is 675 mg, preliminary data from adult studies suggest that even lower DRV doses may be effective, therefore use of 600 mg dose was extended to the entire 25-35 kg weight band.

^dRTV should only be use as a boosting agent in combination with ATV or DRV or to "super boost" LPV/r when given with concomitant rifampicin for TB (see table 5).

^eAt the time of this update, 10 mg and 25 mg DTG film coated tablets were approved for children above 6 years by the FDA (35mg FCT for weight 30 to <40kg, 50mg FCT for weight ≥ 40 kg)⁸ and by the EMA (20mg FTC 15 to < 20, 25mg FCT for 20 to < 30, and 35 FCT for 30 to < 40, 50mg FCT for weight ≥ 40 kg)⁷ based on data from the IMPAACT 1093 trial.⁹ Simplified weight band dosing which deviates from current US FDA and EMA dosing recommendations is being investigated in the Odyssey trial⁵ which supports the use of DTG 50 mg FCT dose for all children ≥ 20 kg. In January 2019 the PAWG reviewed and discussed unpublished data from the Odyssey trial investigating the use of 50 mg FCT in children weighting 20-25 kg¹⁰. The PAWG members acknowledged the short follow up and limited clinical experience, but had no major concerns and agreed with recommending the use of 50 mg FCT from 20 kg, as proposed here. Routine standard toxicity monitoring remains of critical importance in light of the current limited experience with this dosing. All children over 20kg receiving 50mg FCT will continue to be followed up in ODYSSEY and toxicity data collected. For adolescents living with HIV weighing more than 30 Kg a fixed dose formulation of TDF 300mg/3TC 300mg/DTG 50mg (TLD) can be used and is preferred.

Table 3 Simplified dosing of child-friendly solid and oral liquid formulations for twice-daily dosing in infants and children 4 weeks of age and older^a

Drug	Strength of paediatric tablets or oral liquid	Number of tablets or MLs by weight-band morning (AM) and evening(PM)										Strength of adult tablet	Number of tablets by weight band 25–34.9 kg		
		3–5.9 kg		6–9.9 kg		10–13.9 kg		14–19.9 kg		20–24.9 kg					
		AM	PM	AM	PM	AM	PM	AM	PM	AM	PM				
Solid formulations															
AZT	Tablet (dispersible) 60 mg	1	1	1.5	1.5	2	2	2.5	2.5	3	3	300 mg	1		
ABC	Tablet (dispersible) 60 mg	1	1	1.5	1.5	2	2	2.5	2.5	3	3	300 mg	1		
NVP ^b	Tablet (dispersible) 50 mg	1	1	1.5	1.5	2	2	2.5	2.5	3	3	200 mg	1		
LPV/r ^c	Tablet 100 mg/25 mg	–	–	–	–	2	1	2	2	2	2	–	3		
	Pellets 40 mg/10 mg	2	2	3	3	4	4	5	5	6	6	–	–		
	Granules 40 mg/10 mg sachet	2	2	3	3	4	4	5	5	6	6	–	–		
DRV ^d	Tablet 75 mg	–	–	–	–	–	–	5	5	5	5	400 mg	1		
RTV ^e	Tablet 25 mg	–	–	–	–	–	–	2	2	2	2	100 mg	1		
	Tablet 50 mg	–	–	–	–	–	–	1	1	1	1	–	–		
RAL ^f	Chewable tablets 25 mg	1	1	2	2	3	3	4	4	6	6	400 mg	1		
	Chewable tablets 100 mg	–	–	–	–	–	–	1	1	1.5	1.5	–	–		

Drug	Strength of paediatric tablets or oral liquid	Number of tablets or MLS by weight-band morning (AM) and evening(PM)								Strength of adult tablet	Number of tablets by weight band
		3-5.9 kg		6-9.9 kg		10-13.9 kg		14-19.9 kg			
		AM	PM	AM	PM	AM	PM	AM	PM	AM	PM
Liquid formulations											
AZT	10 mg/ml	6 ml	6 ml	9 ml	9 ml	12 ml	12 ml	—	—	—	—
ABC	20 mg/ml	3 ml	3 ml	4 ml	4 ml	6 ml	6 ml	—	—	—	—
3TC	10 mg/ml	3 ml	3 ml	4 ml	4 ml	6 ml	6 ml	—	—	—	—
NVP ^b	10 mg/ml	5 ml	5 ml	8 ml	8 ml	10 ml	10 ml	—	—	—	—
LPV/r ^c	80/20 mg/ml	1 ml	1 ml	1.5 ml	1.5 ml	2 ml	2 ml	2.5 ml	2.5 ml	3 ml	3 ml
DRV ^d	100 mg/ml	—	—	—	—	2.5 ml	2.5 ml	3.5 ml	3.5 ml	—	—
RTV	80 mg/ml	—	—	—	—	0.5 ml	0.5 ml	0.6 ml	0.6 ml	—	—
RAL ^e	10 mg/ml (Oral granules for suspension: 100 mg/sachet)	3 mL	3 mL	5 mL	5 mL	8 mL	8 mL	10 mL	10 mL	—	—

^a See table 4 for dosing recommendations for infants younger than 4 weeks old. Doses for this age group are reduced to account for the decreased ability to excrete and metabolize medications. For infants who are at least 4 weeks of age but less than 3 kg, immaturity of renal and hepatic pathways of elimination are less of a concern but uncertainty still exists on the dosing of ARVs in preterm and low birth weight infants.

^b NVP dose escalation with half dose for 2 weeks when initiating ART is still recommended to avoid toxicity from high initial NVP levels. However, secondary analysis from the (CHAPAS)-1 trial suggested that younger children have a lower risk of toxicity, and consideration can be given to starting with a full dose¹. Please note that this regimen is no longer recommended and should only be used in special circumstances where other age appropriate formulations are not available.

^c LPV/r liquid requires a cold chain during transport and storage. The LPV/r heat-stable formulation must be swallowed whole and should not be split, chewed, dissolved or crushed. Adult 200/50 tablet could be used for patients 14-24.9kg (1 tab am and 1 tab pm), and for patients 25-34.9kg (2 tab am and 1 tab pm). LPVr pellets formulation should not be used in infants younger than 3 months. More details on the administration of LPVr pellets can be found at <https://www.who.int/hiv/pub/toolkits/lat-factsheet-lopinavir-ritonavir/en/>. This dosing schedule applies to equivalent solid dosage forms such as LPV granules which can be use from 2 weeks of age. Since supply is currently constrained both pellets and granules should be discouraged for children above 14 kg who should receive LPV/r 100/25mg tablets instead. Info on LPVr formulations for children available at: <https://www.anprocurementworkinggroup.org/lpv-r-supply>

^d DRV, to be used in children older than 3 years, must be administered with 0.5 ml of RTV 80 mg/ml oral suspension if less than 15 kg and with RTV 50mg using 25mg or 50 mg solid formulation) in children 15 to 30 kg. RTV 100 mg tablets can be used as booster if lower-strength RTV tablets are not available. This is based on limited experience suggesting good acceptability and tolerability. But no efficacy data.

^e RTV should only be used at this dose as a boosting agent in combination with ATV or DRV.

^f RAL granules are approved from birth. Feasibility and acceptability of such formulations has not been widely investigated and concerns have been raised regarding administration in resource limited settings. Due to the administration challenges presented by the granule formulation the use of the 25 mg chewable tablets as dispersible has been endorsed by the PAWG for infants and children older than 4 weeks and weighing at least 3 kg. This was largely based on *in vitro* data on solubility and bioequivalence between tablets and granules⁶ as well as considering the limited availability of adequate alternatives for this age group. However, findings from a feasibility and acceptability assessment conducted in South Africa demonstrate that administration of RAL granules in rural settings is feasible as long as supported with adequate training and counselling.

Table 4 Drug dosing of liquid formulations in infants less than 4 weeks of age^a

Drug	Strength of oral liquid	2-3 kg		3-4 kg		4-5 kg	
		AM	PM	AM	PM	AM	PM
AZT	10 mg/mL	1 mL	1 mL	1.5 mL	1.5 mL	2 mL	2 mL
NVP	10 mg/mL	1.5 mL	1.5 mL	2 mL	2 mL	3 mL	3 mL
3TC	10 mg/mL	0.5 mL	0.5 mL	0.8 mL	0.8 mL	1 mL	1 mL
LPV/r ^b	80/200 mg/mL	0.6 mL	0.6 mL	0.8 mL	0.8 mL	1 mL	1 mL
RAL	Granules 40 mg/10 mg sachet (Oral granules for suspension: 100 mg/sachet) ^c	<1 week >1 week	— 0.4 mL (once daily) ^c	— 0.5 mL (once daily) ^c	2 2	2 0.7 mL (once daily) ^c	2 1.5 mL 1.5 mL

^a PK data in preterm infants are available only for AZT; there is limited data and considerable uncertainty of appropriate dosing for NVP, RAL and 3TC in preterm and low birth weight infants. In addition, LPV/r solution should not be given to preterm infants until they have reached ≥2 weeks gestational age, because of the risk of adverse effects that may occur in this population. This guidance will be updated when more evidence is available from ongoing trials.

^b Do not use LPV/r solution in infants aged >2 weeks of age. LPV pellets should not be used in infants younger than 3 months. More details on the administration of LPV pellets can be found at <https://www.who.int/hiv/pub/toolkits/atr-factsheet-lopinavir-ritonavir/en/>. Due to lack of clinical data to fully inform the use of LPVr granules in neonates, these dosing recommendations were developed on the basis of the current US FDA approval (supporting use of LPVr granules from 2 weeks) and considering the substantial uncertainty which exist particularly for neonates weighing 2-3 kg. Where no other formulation exist, 1 sachet twice a day could be considered for neonates above 2 weeks who are 2-3 kg in order to minimize the risk of potential toxicity with overdosing.

^c RAL granules for oral suspension should use in neonates of at least 2 kg and be administered in once a day during the first week of life (http://www.merck.com/product/usa/pi_circulars/itsentress/itsentress_pi.pdf) and twice a day afterwards.

Table 5 Dosing for RTV super-boosting of LPV/r for children receiving rifampicin-containing TB treatment^a

Drug	Strength of paediatric tablets or oral liquid	Number of tablets or MLs by weight-band morning (AM) and evening (PM)										Strength of adult tablet 3–5.9 kg 6–9.9 kg 10–13.9 kg 14–19.9 kg 20–24.9 kg 25–34.9 kg	Number of tablets by weight band AM PM AM PM AM PM AM PM AM PM		
		For children able to swallow tablets					For children unable to swallow tablets								
LPV/r ^b	Tablet 100/25 mg	—	—	—	—	—	2	1	2	2	2	100/25 mg	3		
RTV	Tablet 100 mg	—	—	—	—	—	1	1	1	2	1	2			
	Tablet 50 mg	—	—	—	—	—	2	2	3	3	3	3			
	Tablet 25 mg	—	—	—	—	—	4	4	6	6	6	6			
LPV/r	Oral solution ^c 80/20 mg/ml	1 ml	1 ml	1.5 ml	1.5 ml	2 ml	2.5 ml	2.5 ml	3 ml	3 ml	—	—	—		
	Pellets ^d 40 mg/10 mg	2	2	3	3	4	4	5	5	6	6	—	—		
	Granules 40 mg/10 mg sachet	2	2	3	3	4	4	5	5	6	6	—	—		
RTV ^e	Oral solution 80 mg/ml	0.8 ml	0.8 ml	1.2 ml	1.2 ml	1.5 ml	2 ml	2 ml	2.3 ml	2.3 ml	—	—	—		
	Powder 100 mg/packet	—	—	1	1	1	1	1	2	1	2	—	—		

^a Suggested RTV dose for super-boosting to achieve the same dose as LPV in mg, in a ratio equal or approaching to 1:1. This dosing approach is supported by a study which explored this approach in young children receiving LPV/r¹².

^b The LPV/r heat-stable tablet formulation must be swallowed whole and should not be split, chewed, dissolved or crushed. Adult 200/50 tablet could be used for patients 14–24.9 kg (1 tab am and 1 tab pm) and for patients 25–34.9 kg (2 tab am and 1 tab pm).

^c LPV/r liquid requires a cold chain during transport and storage.

^d LPV/r pellets formulation should not be used in infants younger than 3 months. More details on the administration of LPV/r pellets can be found at <https://www.who.int/hiv/pub/toolkits/arts-factsheet/lopinavir-ritonavir/en/>. The dosing schedule provided applies to equivalent solid dosage forms that may become available such as LPV/r granules, which are approved by US FDA for use from 2 weeks of life.

^e RTV oral solution dosing is based on the dosing tested in the trial that supports the use of super boosting.

Table 6 Simplified dosing of isoniazid (INH) and co-trimoxazole (CTX) prophylaxis for infants and children who are at least 4 weeks of age

Drug	Strength of paediatric tablet or oral liquid	Number of tablets or ml by weight band once daily			Strength of adult tablet	Number of tablets by weight band
INH		3–5.9 kg	6–9.9 kg	10–13.9 kg	14–19.9 kg	20–24.9 kg 25–34.9 kg
	100 mg	0.5	1	1.5	2	2.5
CTX (SMX/TMP)	Suspension 200/40 per 5 ml	2.5 ml	5 ml	5 ml	10 ml	300 mg –
	Tablets (dispersible) 100/20 mg	1	2	2	4	–
	Tablets (scored) 400/80 mg	–	0.5	0.5	1	400 mg/80 mg 2
	Tablets (scored) 800/160 mg	–	–	–	0.5	800 mg/160 mg 1
	INH/(SMP/ TMP)/B6 ^a	Tablets (scored) 300 mg/ (800mg/160mg)/25 mg	–	–	0.5	300mg/ (800mg/160mg)/25mg 1

^a A scored tablet (150mg/(400mg/80mg)/12.5 mg) is under development.

Optimal Paediatric ARV Formulary

In recent years, a number of improved ARV formulations have become available, such as dispersible, scored FDC tablets which has replaced use of traditional liquid formulations. These products have greatly simplified the delivery of paediatric HIV treatment in low-income settings; however, the proliferation of options, has resulted in a multiplicity of formulations across regimens and weight- bands. Economies of scale are used by generic manufacturers to maintain affordable pricing but fragmentation of demand across too many duplicative products creates instability in the reliable supply of paediatric ARV dosage forms and complicates procurement and supply chain management.

Partners of the ARV procurement working group (APWG)¹³ and of the Global Accelerator for paediatric formulations (GAP-f)¹⁴ provide formulary guidance to programmes on selection of optimal paediatric ARVs which have been defined using a robust set of criteria. The formulary was first developed in 2011 but is routinely revised to correspond to current WHO guidelines and available products. The current Optimal formulary was last revised in June 2018 and is currently a list of 8 products that deliver recommended and appropriate first and second line regimens across all paediatric weight bands. Programs are encouraged to procure paediatric dosage forms that are included on the Optimal Paediatric ARV Formulary. During periods of transitions or in special circumstances (eg. Neonatal treatment, TB co treatment and third line), dosage forms included on the ARV Limited-use formulary provide appropriate coverage¹⁵.

The need for new formulations

As part of the GAP-f, the work of the Paediatric Antiretroviral Working Group (PAWG) and the Paediatric ARV Drug Optimisation (PADO)^{16,17,18} groups continue to highlight the urgent need for better age appropriate formulations for infants and children living with HIV. A number of solid formulations are under final stage of development (ABC/3TC/LPV/r granules as well as ABC/3TC/EFV and DTG 10 mg scored dispersible tablets). In addition, the availability of co-formulated DRV/r in a heat-stable fixed-dose combination is critical to facilitate treatment sequencing and uptake of future paediatric 2nd and 3rd line regimens. A number of formulations containing approved ARVs for paediatric use have been formally prioritised and are listed in table 6. Finally, additional formulations containing newer drugs for which there is currently no paediatric indication were considered and the central future role of DTG and TAF in optimizing dose, sequencing and harmonization across age groups was highlighted.

In moving towards promoting drug optimisation for children and adolescents, WHO will continue to work to simplify prescribing, dispensing and dosing guidance and work with the pharmaceutical industry (originator and generic) and other partners to develop more practical recommendations on the range of formulations required to safely accelerate the scaling up of ART for children.

Box 1 Anticipated simplified dosing for formulations under development

Drug	Strength of dosage form (mg)	Number of tablets or sprinkle capsules or sachets by weight band									
		3–5.9 kg		6–9.9 kg		10–13.9 kg		14–19.9 kg		20–24.9 kg	
		AM	PM	AM	PM	AM	PM	AM	PM	AM	PM
ABC/3TC/PI/r	30 mg/15 mg/40 mg/10 mg granules	2	2	3	3	4	4	5	5	6	6
DRV/r	120 mg/20 mg tablet	—	—	—	—	2	2	3	3	3	—
ABC/3TC/EFV	150 mg/75 mg/150 mg tablet	—	—	—	—	1.5	—	2	2.5	3	4
DTG ^a	Scored dispersible 10 mg tablet	1	—	—	—	1.5	—	2	2.5	3	4
ABC/3TC/DTG ^b	Dispersible 60 mg/30 mg/5 mg tablet	2	—	3	—	4	—	5	6	—	—

^a This dosing was confirmed by the PAWG in May 2019 on best available information. However, optimal dosing of DTG in children below 20 kg is still being investigated and this proposed dosing will be revised as soon as more evidence is gathered from ongoing studies.

^b This dosage form is the one identified by the PADO4 group¹⁸ as the most likely to deliver appropriate dose based on the best available information. However, the group strongly emphasized the importance of validating this dosing and ratio as soon as final dosing for DTG is approved down to 4 weeks of age. Scored tablets with doubled-strength of each component drug (ABC/3TC/DTG 120/60/10 mg) would enable reduction of the pill burden for children, but difficulties with assuring accuracy of dosing when scoring a triple-drug.

- 1 PAWG members: Elaine Abrams (ICAP at Columbia University, USA); Mo Archary (University of KwaZulu-Natal, South Africa); Yodit Belew (US Food and drug administration HHS, USA); Brookie Best (University of San Diego, USA); David Burger (Radboud University Nijmegen Medical Centre, The Netherlands); Jessica Burry (Access Campaign MSF, Switzerland); Edmund Capparelli (University of California San Diego, USA); Deborah Carpenter (Center for Disease Control, USA); Polly Clayden (HIV i-Base, UK); Diana Clarke (Boston Medical Center, USA); Timothy R. Cressey (Program for HIV Prevention and Treatment-IRD/Harvard T.H Chan School of Public Health & Chang Mai University, Thailand); Angela Colbers (Radboud University Nijmegen Medical Centre, The Netherlands); Mutsa Dangarembizi (University of Zimbabwe, Zimbabwe); Paolo Denti (University of Cape Town, South Africa); Andrea Ecker (European Medicines Agency, The Netherlands); Shaffiq Essajee (UNICEF, USA); Carlo Giaquinto (University of Padova, Italy); Diana Gibb (MRC Clinical Trials Unit, United Kingdom); Rohan Hazra (National Institute of Child Health and Human Development, USA); Maria Kim (Baylor College of Medicines, Malawi); Marc Lallement (PHPT Foundation, Thailand); Janice Lee (Drugs for Neglected Diseases initiative, Switzerland); Linda Lewis (Clinton Health Access Initiative, USA); Shahin Lockman (Harvard T.H Chan School of Public Health, USA); Mark H. Mirochnick (Boston Medical Center, USA); Lynne Mofenson (Elizabeth Glaser Paediatric AIDS Foundation, USA); Victor Musiime (Joint Clinical Research Center, Uganda); Elizabeth Obimbo (University of Nairobi, Kenya); Atieno Ojoo (UNICEF, Denmark); Fernando Pascual (Medicines Patent Pool, Switzerland); Jorge Pinto (Federal University of Minas Gerais Belo Horizonte, Brazil); Natella Rakhmanina (Elizabeth Glazer Paediatric AIDS Foundation, USA); Pablo Rojo (Hospital de 12 Octubre Madrid, Spain); Ted Ruel (University of California San Francisco, USA); George Siberry (USAID, USA); Nandita Sugandhi (ICAP at Columbia University, USA); Marissa Vicari (International AIDS Society, Switzerland); Melynda Watkins (Clinton Health Access Initiative, USA); Hilary Wolf ((Office of the U.S. Global AIDS Coordinator U.S. Department of State, USA)).
- 2 http://www.ema.europa.eu/docs/en_GB/document_library/EPAR_-_Product_Information/human/002753/WC500160680.pdf
- 3 https://www.accessdata.fda.gov/drugsatfda_docs/label/2017/204790s014lbl.pdf
- 4 <https://clinicaltrials.gov/ct2/show/NCT01302847?term=P1093&rank=2>
- 5 Turkova et al .2018 10th International Workshop on HIV Paediatrics 20 – 21 July 2018, Amsterdam, The Netherlands
- 6 Teppier H, Thompson K, Chain A, Mathe M, Nachman S, Clarke D. Crushing of Raltegravir (RAL) chewable tablets for administration in infants and young children. Abstract 37. 9th International Workshop on HIV Paediatrics 21 – 22 July 2017, Paris, France
- 7 https://www.accessdata.fda.gov/drugsatfda_docs/label/2017/207561s013lbl.pdf
- 8 <https://www.ncbi.nlm.nih.gov/pubmed/29206747>
- 9 <https://clinicaltrials.gov/ct2/show/NCT02259127?term=DOLUTEGRAVIR+CHILDREN&cond=HIV&rank=3>
- 10 Bollen et al. <http://www.croiconference.org/sessions/adult-dolutegravir-50mg-tablets-children-living-hiv-weighing-20>
- 11 <https://www.ncbi.nlm.nih.gov/pubmed/22739394>
- 12 Rabie H, Denti P, Lee J, Masango M, Coovadia A, Pillay S, Liberty A, McIlheron H, Cotton M, Lallement M. Lopinavir/Ritonavir 1:1 super-boosting overcomes rifampicin interactions in children. Proceedings of the Conference on Retroviruses and Opportunity Infections; 2017 Feb 13-16; Seattle, USA; https://www.dndi.org/wp-content/uploads/2018/04/Rabie_et_al_Super_Boosting_Children_Abstract_CROI2017.pdf
- 13 APWG <https://www.arvprocurementworkinggroup.org/>
- 14 GAPf <http://www.gap-f.org/>
- 15 The 2018 optimal formulary and limited-use list for paediatric ARVs. Policy brief <http://www.who.int/hiv/pub/paediatric/optimal-paediatric-arv-formulary/en/>
- 16 World Health Organization. Meeting report: Paediatric Antiretroviral Drug Optimization (PADO) Meeting 3. Geneva, Switzerland: WHO; 2016. Available at: <http://www.who.int/hiv/pub/meetingreports/paediatric-arv-optimization-pado3/en/> Accessed on November 29, 2017.
- 17 World Health Organization. Meeting report: Paediatric ARV drug optimization 3 review <http://www.who.int/hiv/pub/meetingreports/pado3-review/en/index1.html>
- 18 <https://www.who.int/hiv/pub/meetingreports/paediatric-arv-optimization-pado4/en/>